

CORONASTEP Report 179 (2024 – Weeks 09 and 10) SARS-CoV-2 Sewage Surveillance in Luxembourg

Summary

After a detailed analysis of the data collected to date, we demonstrated that it was possible to reduce the number of wastewater treatment plants (WWTPs) monitored without affecting the relevance and representativeness of the national data. The 8 WWTPs selected are as follows: Beggen, Schifflange, Bettembourg, Pétange, Mersch, Echternach, Ubersyren and Blesbruck.

From now on, only the results of these eight analyses will be presented in the fortnightly reports.

This report 179 presents the results of SARS-CoV-2 contamination of wastewater at the entrance of the 8 wastewater treatment plants (WWTPs) analysed during weeks 09 and 10 of 2024. At the national level, the SARS-CoV-2 RNA fluxes measured are lower than those measured over the previous two weeks, with a national flux of SARS-CoV-2 around 1.5×10^{11} RNA copies per day per 100,000 population equivalents. Overall, the general trend is still downwards.

Remark: Following the recent adoption of the Act amending the Act of 17 July 2020 on measures to combat the COVID-19 pandemic and the decisions of the Government Council regarding the surveillance of the COVID-19 epidemic, the data on clinical surveillance will no longer be reported on the national and regional charts. Indeed, the number of tests performed has already started to decrease drastically and is therefore no longer as representative of the virus circulation in the population as before.

Table 1 – National level of SARS-CoV-2 contamination of wastewaters in Luxembourg.



Dark green: negative samples for SARS-CoV-2 gene E (-), Green to red: positive samples for SARS-CoV-2 gene E. The intensity of the color is related to the national SARS-CoV-2 flux (RNA copies / day / 100 000 equivalent inhabitants).

2020	
National SARS-CoV-2 Level	Week
Dark Green	Week 3
Dark Green	Week 7
Dark Green	Week 9
Dark Green	Week 11
Light Green	Week 14
Light Green	Week 15
Light Green	Week 16
Light Green	Week 17
Light Green	Week 18
Light Green	Week 19
Light Green	Week 20
Light Green	Week 21
Light Green	Week 22
Light Green	Week 23
Light Green	Week 24
Light Green	Week 25
Light Green	Week 26
Light Green	Week 27
Light Green	Week 28
Light Green	Week 29
Light Green	Week 30
Light Green	Week 31
Light Green	Week 32
Light Green	Week 33
Light Green	Week 34
Light Green	Week 35
Light Green	Week 36
Light Green	Week 37
Light Green	Week 38
Light Green	Week 39
Light Green	Week 40
Light Green	Week 41
Light Green	Week 42
Light Green	Week 43
Light Green	Week 44-1
Light Green	Week 44-2
Light Green	Week 45-1
Light Green	Week 45-2
Light Green	Week 45-3
Light Green	Week 46-1
Light Green	Week 46-2
Light Green	Week 46-3
Light Green	Week 47-1
Light Green	Week 47-2
Light Green	Week 48-1
Light Green	Week 48-2
Light Green	Week 48-3
Light Green	Week 49-1
Light Green	Week 49-2
Light Green	Week 50-1
Light Green	Week 50-2
Light Green	Week 51-1
Light Green	Week 51-2
Light Green	Week 52
Light Green	Week 53


2021	
National SARS-CoV-2 Level	Week
Light Green	Week 01-1
Light Green	Week 01-2
Light Green	Week 02-1
Light Green	Week 02-2
Light Green	Week 03-1
Light Green	Week 03-2
Light Green	Week 04-1
Light Green	Week 04-2
Light Green	Week 05-1
Light Green	Week 06-1
Light Green	Week 06-2
Light Green	Week 07-1
Light Green	Week 07-2
Light Green	Week 08-1
Light Green	Week 08-2
Light Green	Week 09-1
Light Green	Week 09-2
Light Green	Week 10-1
Light Green	Week 10-2
Light Green	Week 11-1
Light Green	Week 11-2
Light Green	Week 12-1
Light Green	Week 12-2
Light Green	Week 13-1
Light Green	Week 13-2
Light Green	Week 14-1
Light Green	Week 14-2
Light Green	Week 15-1
Light Green	Week 15-2
Light Green	Week 16-1
Light Green	Week 16-2
Light Green	Week 17-1
Light Green	Week 17-2
Light Green	Week 18-1
Light Green	Week 18-2
Light Green	Week 19
Light Green	Week 20-1
Light Green	Week 20-2
Light Green	Week 21
Light Green	Week 22-1
Light Green	Week 22-2
Light Green	Week 23-1
Light Green	Week 23-2
Light Green	Week 24-1
Light Green	Week 24-2
Light Green	Week 25
Light Green	Week 26-1
Light Green	Week 26-2
Light Green	Week 27-1
Light Green	Week 27-2
Light Green	Week 28-1
Light Green	Week 28-2
Light Green	Week 29-1
Light Green	Week 29-2
Light Green	Week 30-1
Light Green	Week 30-2

2021		2022	
National SARS-CoV-2 Level	Week	National SARS-CoV-2 Level	Week
Light Green	Week 31-1	Light Green	Week 01
Light Green	Week 31-2	Light Green	Week 02-1
Light Green	Week 32-1	Light Green	Week 02-2
Light Green	Week 32-2	Light Green	Week 03-1
Light Green	Week 33-1	Light Green	Week 03-2
Light Green	Week 33-2	Light Green	Week 04-1
Light Green	Week 34-1	Light Green	Week 04-2
Light Green	Week 34-2	Light Green	Week 05-1
Light Green	Week 35-1	Light Green	Week 05-2
Light Green	Week 35-2	Light Green	Week 06-1
Light Green	Week 36-1	Light Green	Week 06-2
Light Green	Week 36-2	Light Green	Week 07-1
Light Green	Week 37-1	Light Green	Week 07-2
Light Green	Week 37-2	Light Green	Week 08-1
Light Green	Week 38-1	Light Green	Week 08-2
Light Green	Week 38-2	Light Green	Week 09-1
Light Green	Week 39-1	Light Green	Week 09-2
Light Green	Week 39-2	Light Green	Week 10-1
Light Green	Week 40-1	Light Green	Week 10-2
Light Green	Week 40-2	Light Green	Week 11-1
Light Green	Week 41-1	Light Green	Week 11-2
Light Green	Week 41-2	Light Green	Week 12-1
Light Green	Week 42-1	Light Green	Week 12-2
Light Green	Week 42-2	Light Green	Week 13-1
Light Green	Week 43-1	Light Green	Week 13-2
Light Green	Week 43-2	Light Green	Week 14-1
Light Green	Week 44-1	Light Green	Week 14-2
Light Green	Week 44-2	Light Green	Week 15-1
Light Green	Week 45-1	Light Green	Week 15-2
Light Green	Week 45-2	Light Green	Week 16-1
Light Green	Week 46-1	Light Green	Week 16-2
Light Green	Week 46-2	Light Green	Week 17-1
Light Green	Week 47-1	Light Green	Week 17-2
Light Green	Week 47-2	Light Green	Week 18-1
Light Green	Week 48-1	Light Green	Week 18-2
Light Green	Week 48-2	Light Green	Week 19-1
Light Green	Week 49-1	Light Green	Week 19-2
Light Green	Week 49-2	Light Green	Week 20-1
Light Green	Week 50-1	Light Green	Week 20-2
Light Green	Week 50-2	Light Green	Week 21-1
Light Green	Week 51-1	Light Green	Week 21-2
Light Green	Week 51-2	Light Green	Week 22-1
Light Green	Week 01	Light Green	Week 22-2
Light Green	Week 02-1	Light Green	Week 23-1
Light Green	Week 02-2	Light Green	Week 23-2
Light Green	Week 03-1	Light Green	Week 24-1
Light Green	Week 03-2	Light Green	Week 24-2
Light Green	Week 04-1	Light Green	Week 25-1
Light Green	Week 04-2	Light Green	Week 25-2
Light Green	Week 05-1	Light Green	Week 26-1
Light Green	Week 05-2	Light Green	Week 26-2
Light Green	Week 06-1	Light Green	Week 27
Light Green	Week 06-2	Light Green	Week 28
Light Green	Week 07-1	Light Green	Week 29
Light Green	Week 07-2	Light Green	Week 30

2022	
National SARS-CoV-2 Level	Week
Light Green	Week 08-1
Light Green	Week 08-2
Light Green	Week 09-1
Light Green	Week 09-2
Light Green	Week 10-1
Light Green	Week 10-2
Light Green	Week 11-1
Light Green	Week 11-2
Light Green	Week 12-1
Light Green	Week 12-2
Light Green	Week 13-1
Light Green	Week 13-2
Light Green	Week 14-1
Light Green	Week 14-2
Light Green	Week 15-1
Light Green	Week 15-2
Light Green	Week 16-1
Light Green	Week 16-2
Light Green	Week 17-1
Light Green	Week 17-2
Light Green	Week 18-1
Light Green	Week 18-2
Light Green	Week 19-1
Light Green	Week 19-2
Light Green	Week 20-1
Light Green	Week 20-2
Light Green	Week 21-1
Light Green	Week 21-2
Light Green	Week 22-1
Light Green	Week 22-2
Light Green	Week 23-1
Light Green	Week 23-2
Light Green	Week 24-1
Light Green	Week 24-2
Light Green	Week 25-1
Light Green	Week 25-2
Light Green	Week 26-1
Light Green	Week 26-2
Light Green	Week 27
Light Green	Week 28
Light Green	Week 29
Light Green	Week 30
Light Green	Week 31
Light Green	Week 32
Light Green	Week 33
Light Green	Week 34
Light Green	Week 35
Light Green	Week 36
Light Green	Week 37
Light Green	Week 38
Light Green	Week 39
Light Green	Week 40
Light Green	Week 41
Light Green	Week 42
Light Green	Week 43

Figure 1 – RT-qPCR quantification time-course monitoring of SARS-CoV-2 (*E* gene) in Luxembourgish wastewater samples from December 2019 to March 2024. Grey squares: daily confirmed cases for Luxembourgish residents (<https://data.public.lu/fr/datasets/donnees-covid19/>), Blue dots: cumulative SARS-CoV-2 flux (RNA copies / day / 100 000 equivalent inhabitants).

a) Linear scale



b) Log_{10} scale







Figure 2 – RT-qPCR quantification time-course monitoring of SARS-CoV-2 (E gene) in the eight selected wastewater treatment plants from March 2020 to March 2024. Grey squares: daily confirmed cases for the contributory area of each wastewater treatment plant, colored dots: SARS-CoV-2 flux (RNA copies / day / 10,000 equivalent inhabitants).




Pétange




Bettembourg




Bleesbruck



Mersch



Uebersyren



Echternach




Table 3- Sewage sampling since the beginning of the CORONASTEP study

WWTP	2019	2020	2021	2022	2023	2024	Total
Beggen		52	92	77	50	10	281
Bettembourg		45	91	75	50	10	271
Schifflange	4	56	96	76	51	10	293
Bleesbrück		44	92	77	51	10	274
Mersch		47	91	77	51	10	276
Pétange	4	51	96	77	51	10	289
Hesperange		35	48	50	51	3	187
Echternach		36	79	77	50	9	251
Uebersyren		38	78	77	50	9	252
Grevenmacher		38	81	76	51	2	248
Troisvierges		38	92	77	51	3	261
Boevange sur Attert		20	48	51	51	3	173
Wiltz		28	92	77	51	3	251
Total	8	528	1076	944	659	92	3307

Materials and Methods

Sewage samples

From March 2020 to March 2024, up to thirteen wastewater treatment plants (WWTPs) were sampled at their inlet according to the planning presented in Table 3. Since January 2024, the number of WWTPs followed was reduced to eight. The operators of the WWTPs collected a 24-hour composite sample according to their routine sampling procedure. Composite samples were stored at 4°C until sample processing.

Sample processing

The samples were transported to the laboratory at 4°C and viral RNA was isolated on sampling day. Larger particles (debris, bacteria) were removed from the samples by centrifugation at 2,400 x g for 20 min at 4°C. A 120 mL of supernatant was filtered through Amicon® Plus-15 centrifugal ultrafilter with a cut-off of 10 kDa (Millipore) by centrifugation at 3,220 x g for 25 min at 4°C. The resulting concentrate was collected and 140 µL of each concentrate was then processed to extract viral RNA using the QIAamp Viral RNA mini kit (Qiagen) according to the manufacturer's protocol. Elution of RNA was done in 60 µL of elution buffer.

Real-time One-Step RT-PCR

Samples were screened for the presence of *Sarbecovirus* (*Coronaviridae*, *Betacoronaviruses*) and/or SARS-CoV-2 virus RNA by two distinct real-time one-step RT-PCR assays, targeting the E gene (Envelope small membrane protein) and the N gene (nucleoprotein). The E gene real-time RT-PCR can detect *Sarbecoviruses*, i.e. SARS-CoV, SARS-CoV-2 and closely related bat viruses. In the context of the COVID19 pandemic, it can be assumed that only SARS-CoV-2 strains will be detected by this assay given that SARS-CoV virus has been eradicated and other bat viruses do not commonly circulate in the human population. The E gene assay is adapted from Corman et al. [17]. The N gene real-time RT-PCR assay (N1 assay) specifically detects SARS-CoV-2 virus. It is adapted from the CDC protocol¹. The two primers/probe sets are presented in Table 3. The RT-qPCR protocols and reagents were all provided by the LIH.

Table 4 – RT-qPCR primer-probe sets.

Target	Primer name	Primer sequence (5' to 3')	References
E gene	E_Sarbeco_F1	5-ACAGGTACGTTAATAGTTAATAGCGT-3	Corman et al., 2020
	E_Sarbeco_R2	5-ATATTGCAGCAGTACGCACACA-3	
	E_Sarbeco_P1	5'-FAM-ACACTAGCCATCCTTACTGCGCTTCG-BHQ1	
N gene	2019-nCoV_N1_Fw	5'-GAC CCC AAA ATC AGC GAA AT-3'	CDC, 2019
	2019-nCoV_N1_Rv	5'-TCT GGT TAC TGC CAG TTG AAT CTG-3'	
	2019-nCoV_N1 Probe	5'-FAM-ACC CCG CAT TAC GTT TGG TGG ACC-BHQ1-3'	

Each reaction contained 5 µL of RNA template, 5 µL of TaqPath 1-step RT-qPCR MasterMix (A15299, Life Technologies), 0.5 µL of each primer (20 µM) and probe (5 µM) and the reaction volume was adjusted to a final volume of 20 µL with molecular biology grade water. Thermal cycling reactions were carried out at 50 °C for 15 min, followed by 95 °C for 2 min and 45 cycles of 95 °C for 3 sec and 58°C (E gene) or 55°C (N gene) for 30 sec using a ViiA7 Real-Time PCR Detection System (Life Technologies). Reactions were considered positive (limit of detection – LOD) if the cycle threshold (Ct value) was below 40 cycles.

¹ <https://www.cdc.gov/coronavirus/2019-ncov/downloads/rt-pcr-panel-primer-probes.pdf>

Controls

A non-target RNA fragment commercially available (VetMAX™ Xeno™ IPC and VetMAX™ Xeno™ IPC Assay, ThermoFischer Scientific) was added to the viral RNA extract from sewage concentrates as an internal positive control (IPC). This IPC-RNA is used to control the performance of the RT-qPCR (E gene) and to detect the presence of RT-qPCR inhibitors.

Viral RNA copies quantification of both targeting genes in wastewater samples was performed using RT-qPCR standard curves generated using EDX SARS-CoV-2 Standard (Biorad). This standard is manufactured with synthetic RNA transcripts containing 5 targets (E, N, S, ORF1a, and RdRP genes of SARS-CoV-2, 200,000 copies/mL each). Using such a standard, the limits of quantification (LOQ) of both RT-qPCR assays were estimated to 1 RNA copy per reaction (Figure 6).




Figure 6 – RT-qPCR standard curves established for both target genes (E gene and N gene) of SARS-CoV-2 using a commercially available standard (Biorad).


Data interpretation

A sample is declared positive for the presence of SARS-CoV-2 if both targets (E and N gene) are detected with Ct values less than or equal to the LOQ. If only one target is detected or if target genes are detected with Ct values between the LOD and the LOQ, samples are reported as presumptive positive (+/-). A sample is declared negative when no target genes are detected (Ct values superior to the LOD).

In case of presumptive positive, sample is tested again using another RT-qPCR detection assay (Allplex 2019-nCoV Assay, Seegene). This commercially available detection kit is a multiplex real-time RT-PCR assay for simultaneous detection of three target genes of SARS-CoV-2 in a single tube. The assay is designed to detect RdRP and N genes specific for SARS-CoV-2, and E gene specific for all *Sarbecovirus* including SARS-CoV-2.

As shown in Figure 7, a highly significant correlation (Pearson Correlation, $R^2=0.964$, $p = 5.979 \cdot 10^{-24}$) was obtained between the SARS-CoV-2 RNA concentrations estimated using the E gene and the N gene, respectively. Therefore, only the E gene results were presented in this report.

Figure 7 - Relationship between the SARS-CoV-2 RNA concentration (RNA copies / L of wastewater) estimated by the both distinct RT-qPCR systems targeting the E and N gene, respectively (n=415),



Acknowledgments

This work is supported by the Fond National de la Recherche (FNR) under project 14806023 - CORONASTEP+ and is conducted in collaboration with the Luxembourg Institute of Health (LIH), the “Laboratoire National de Santé” (LNS) and the University of Luxembourg (LCSB).

In addition, the authors of this report would like to thank all the wastewater syndicates (SIACH, SIVEC, STEP, SIDERO, SIDEN and SIDEST), the “Ville du Luxembourg”, the Hesperange city as well as the “Administration de la Gestion de l’Eau” (AGE) for their kind and valuable assistance in the sample collection, the acquisition of wastewater parameters and the collection of demographic data. The authors would also like to thank the Ministry of Health and the Inspection Sanitaire for their valuable contribution in providing the COVID-19 data at the national and regional scale.