
RESEARCH GROUP

Microbiome Engineering



The Microbiome Engineering group works at the interface of engineering, microbiology, and computational biology to develop bio-based processes to produce bioenergy and biomolecules from biomass. The group is especially interested in the process of anaerobic digestion and biomethanation, studying both natural and engineered systems. In addition, the group has been applying its internally optimized molecular biology tools to study, in a high-throughput manner, microbiomes from lab-scale reactors and full-scale energy units, as well as communities in soil, animal guts, cave deposits, etc.

Main expertise fields

- Anaerobic digestion (AD) is the most advanced technology to convert carbon-rich biomass, including lignocellulosic waste, into energy (methane-rich biogas) and value-added products, such as biofertilizers. Methane-rich biogas can be further upgraded to natural gas quality, via a process called biomethanation.
 Lignocellulose degradation. Lignocellulose is the major structural component of plant biomass-rich habitats, with the objective of further valorizing their lignocellulosity capacities for industrial applications.
 Soil Microbiome. The soil microbiome regroups mainly bacteria, archaea, viruses, fungi, protists, and other small eukaryotes such as diatoms. Soil microorganisms play key roles in carbon and nutrient cycling by decomposing soil organic matter and transforming important nutrients for soil fertility, thus
- determining the productivity of agroecosystems. In addition, despite their importance in food security and climate change, most of the soil microbes remain largely uncharacterized.

 Wastewater treatment and pollutant removal, focusing on a biological process (e.g. aerobic granular sludge reactors) or the removal of recalcitrant compounds via adsorption or membrane operations

Research challenges

- draw and a state of the group develops innovative reactor designs and process monitoring tools to boost the process of biomass (organic waste, including lignocellulose) and gaseous waste valorization, contributing to the development of the biorefinery sector, as explained more

- To characterise the microbiome, the group relies on high-throughput sequencing, employing metagenomics and metatranscriptomics, as well as metaproteomics and metabolomics. By doing this, the group has recently shown that propionate oxidizing bacteria of the candidate phylum Cloacimonetes, can largely improve the resilience of the anaerobic digestion process, greatly boosting methane production from carbon-rich waste, at high organic loading rates. Currently, the group exploits their potential as a healthy implant of the anaerobic digestion reactors.

 The group has prosposed exploiting the potential of multivariate statistical process monitoring on biogas composition to detect free ammonia intoxication at an early stage in anaerobic digestion reactors;

 Recently, the group has developed a dense membrane gas capture reactor to promote efficient biomethanation under mesophilic conditions. Currently a pilot-scale prototype has been implemented on a farm and tested under real-scale conditions. Preliminary lab-scale experiments have proven high rates of CO2 conversion to methane, with the methane content in the biogas > 95% (i.e. natural gas quality biogas).

The Microbiome Engineering has investigated the highly specialized higher termite gut system, anaerobic digestion microbiome and soil. The most interesting outcomes include the following:

- For the first time, the group has identified and characterized multifunctional cellulases from the termite gut microbes, which are capable of simultaneously cleaving linkages between cellulose, xylans and mannans moieties.
 The application of an integrated omics approach to the anaerobic digestion microbiome allowed the group to identify members of the phylum Planctomycetes as previously unexplored robust biomass degrading bacteria, of high industrial relevance.

One of the group's missions is to promote anaerobic digestion residue as an environmentally benign biofertilizer which could contribute to the mitigation of nitrate leaching in agricultural soils and promote carbon sequestration in grasslands. The most interesting outcomes are detailed below

- The diversity of bacterial communities is pH dependent, however, arable soils have more diverse microbiomes than grasslands and forests.
 Granulated biogas residue is a biofertilizer with the potential to supply nutrients to soil biota over time, and promote carbon sequestration in grassland soils, and thereby advance agricultural sustainability while contributing to climate change mitigation.
 The (partial) substitution of chemical fertilizers by raw digestate reduces the concentration of intracts in the soil without having a negative impact on the yield and N content of the biomass.

- Characterizations of microbial communities (microbiomes and vironnes whole viral communities) and as

 Isolation and characterization of aerobic and anaerobic bacteria of industrial relevance.
 Discovery, production and characterization of arcarbohydrate active enzymes for tailored solutions.

Main assets

- Lab-scale and pilot-scale anaerobic digestion reactors, including 200 continuously stirred-tank reactors (CSTRs) with 2L capacity; four CSTRs with 100L capacity; eight anaerobic baffled reactors (ABRs) with 2L capacity and four compartmented ABRs with 100 L capacity.

- Two in-house servers and easy access to the LIST HPC facilities and other HPC resources available in Luxembourg.
 Anaerobic and microaerophilic cultivation chambers for the isolation and growing of microorganisms. The complete cultivation infrastructure includes several hundreds of sealed cultivation bottles, and instrumentation for anaerobic media generation, incubators, shakers, etc.

- Westerholm, M., Calusinska, M. and Dolfing, J. <u>Syntrophic propionate oxidizing bacteria in methanogenic systems</u> (2022) FEMS Microbiology Reviews, Volume 46, Issue 2, fuab057
 Calusinska, M., Marynowska, M., Bertucci, M. et al. <u>Integrative omics analysis of the termite gut system adaptation to Miscarthus diet identifies lignocellulose degradation enzymes (2020) Commun Biol 3, 275
 Calusinska, M., Goux, X., Forsepre, M. Muller EEL Wilmses P and Deficioses P. A year of monitoring 20 mesophilic full-scale biorgeactors reveals the existence of stable but different core microbiomes in bio-waste and wastewater anaerobic digestion systems
 Plattes, M., Goux, X., O'Nagy, O., Untereiner, B., Fernandez Lahore, H.M. Zero-dimensional modelling, and bacterial characterization of an aerobic granular sludge reactor (2023) Journal of Chemical Technology and Biotechnology, 98 (2), pp. 369-380.</u>

Partenaires

Swedish University of Agricultural Sciences, Biorock, Wagner Schaffner, Schroeder & Associés

Contact

5, avenue des Hauts-Fourneaux L-4362 Esch-sur-Alzette tél: +352 275 888 - 1 | LIST.lu

